

Guidelines for Concussion / Mild Traumatic Brain Injury & Persistent Symptoms

Second Edition

For adults (18+ years of age)



Module 10: Persistent Vestibular (Balance / Dizziness) & Vision Dysfunction



Ontario Neurotrauma Foundation
Fondation ontarienne de neurotraumatologie

MODULE 10: PERSISTENT VESTIBULAR (BALANCE/DIZZINESS) & VISION DYSFUNCTION



Ontario Neurotrauma Foundation
Fondation ontarienne de neurotraumatologie

The project team would like to acknowledge the Ontario Neurotrauma Foundation (ONF), who initiated and funded the development of the original guideline, as well as the current update. ONF is an applied health research organization with a focus on improving the quality of lives for people with an acquired brain injury or spinal cord injury, and on preventing neurotrauma injuries from occurring in the first place. ONF uses strategic research funding activity embedded within a knowledge mobilization and implementation framework to build capacity within systems of care. ONF works with numerous stakeholders and partners to achieve its objective of fostering, gathering and using research knowledge to improve care and quality of life for people who have sustained neurotrauma injuries, and to influence policy towards improved systems. The foundation receives its funding from the Ontario Government through the Ministry of Health and Long-Term Care.

Please note, the project team independently managed the development and production of the guideline and, thus, editorial independence is retained.

© Ontario Neurotrauma Foundation 2013

**Ontario Neurotrauma Foundation
90 Eglinton East
Toronto, ON, Canada M4P 2Y3
Tel.: 1 (416) 422-2228
Fax: 1 (416) 422-1240
Email: info@onf.org**

www.onf.org

Published September 2013

The recommendations and resources found within the *Guidelines for Concussion/Mild Traumatic Brain Injury & Persistent Symptoms* are intended to inform and instruct care providers and other stakeholders who deliver services to adults who have sustained or are suspected of having sustained a concussion/mTBI. These guidelines are not intended for use with patients or clients under the age of 18 years. These guidelines are not intended for use by people who have sustained or are suspected of having sustained a concussion/mTBI for any self-diagnosis or treatment. Patients may wish to bring their healthcare and other providers' attention to these guidelines.

The recommendations provided in these guidelines are informed by best available evidence at the time of publication, and relevant evidence published after these guidelines could influence the recommendations made within. Clinicians should also consider their own clinical judgement, patient preferences and contextual factors such as resource availability in clinical decision-making processes.

The developers, contributors and supporting partners shall not be liable for any damages, claims, liabilities, costs or obligations arising from the use or misuse of this material, including loss or damage arising from any claims made by a third party.

Table of Contents

GUIDELINE RECOMMENDATIONS

10. Persistent Vestibular (Balance/Dizziness) & Vision Dysfunction	1
--	---

APPENDICES

10.1: Dix-Hallpike Manoeuvre and Particle Repositioning Manoeuvre (PRM)	5
10.2: Screening Techniques for Vision Dysfunction	6
A: Project Members	7
B: Other Useful Links/References for Resources to Consider	9

TABLES

10.1: Important Components to Include in the Neurological Exam	1
10.2: Post-Trauma Vision Syndrome (PTVS) Definitions	3
10.3: Common Visual Symptoms and Associated Visual Deficits	3

FIGURE

10.1: Clinical Assessment of Balance.....	2
---	---

Unique Features & Symbols in the Current Guideline

Hyperlinks

To improve ease of use, the current guideline has embedded hyperlinks to improve navigation between sections, appendices, and so on. For example, by clicking any heading in the table of contents above, you will be taken directly to that particular section in the current PDF document. Also, anytime there is mention of a particular table, figure, appendix or section, you can simply click on it to go directly to that item.

Symbols



The following symbol has been placed to the left of each guideline recommendation that should be prioritized for implementation. This was determined by expert consensus members during the endorsement/prioritization process, where experts were allowed to provide 20 prioritization votes (see Methodology in the Complete Version). Guideline recommendations with a summed prioritization score greater than 20 are key clinical practice guidelines recommendations for implementation.



The following symbol has been placed to the left of one key guideline recommendation in each of the sections that did not include a recommendation with a prioritization score greater than 20 (determined by expert consensus members during the endorsement/prioritization process).

At the bottom of each page in the current document, there is a hyperlinked footer that can be used to return to the table of contents as desired. Also, clicking “Return to Last Page” will take you back to the previously viewed page. (Note: When scrolling through the pages, the “Return to Last Page” button will only return to the last page that was scrolled through).

10 Persistent Vestibular (Balance/Dizziness) & Vision Dysfunction

Special contributors:
Angela Peddle (vision),
Jennifer Shea (vestibular)

Vestibular (Balance/Dizziness) Dysfunction

Persistent vertigo, dizziness, imbalance, and vision changes are common complaints post mTBI and are often associated with vestibular system impairments.^{1,2} Vestibular deficits can be peripheral in origin, affecting the inner ear, or central, affecting central nervous system integration and output to maintain balance and posture. The peripheral vestibular organs also affect eye movement through the vestibulo-ocular reflex (VOR). Thus, vestibular dysfunction presents as balance impairments and VOR abnormalities.

The most common cause of post-traumatic peripheral vestibular dysfunction is benign paroxysmal positional vertigo (BPPV).³ Patients experience episodes of vertigo, nystagmus, and nausea with sudden changes in position, often including rolling over in bed or looking up. These attacks usually last less than 30 seconds, but can be quite disabling and occur multiple times per day. BPPV is most commonly caused by dislodged otoconia in the posterior semicircular canal (SCC).

Assessment of vestibular function is important following mTBI to identify vestibular deficits, which may benefit from evidence-based interventions. Evaluation should minimally include a balance screen, the Dix-Hallpike manoeuvre and VOR screening. Balance testing should reference normal values to document impairment (see Figure 10.1).^{4,5}

When the history suggests BPPV, posterior SCC involvement can be diagnosed by the Dix-Hallpike manoeuvre (see Appendix 10.1 for more information and Appendix B for links to video demonstrations). VOR abnormalities will often present as nystagmus in one or more directions of gaze. When assessment suggests vestibular dysfunction, vestibular interventions can be considered. Although, historically, medications have been used to suppress vestibular symptoms, including nausea, current evidence does not support this approach. A Cochrane review by Hillier and Hollohan (2007) identifies vestibular rehabilitation (VR) as an effective intervention for unilateral peripheral vestibular dysfunction.¹ Weaker evidence also suggests VR may be helpful for central vestibular dysfunction.⁶ VR is typically provided by a specialized physiotherapist and involves various movement-based regimens to bring on vestibular symptoms and desensitize the vestibular system, coordinate eye and head movements, and improve functional balance and mobility. However, for the specific treatment of BPPV, Hillier and Hollohan (2007) conclude that canalith or particle repositioning manoeuvres are more effective than VR techniques.¹

Table 10.1 Important Components to Include in the Neurological Exam

Vision	<ul style="list-style-type: none"> • Acuity • Tracking • Saccades • Nystagmus • Vergence
Auditory	<ul style="list-style-type: none"> • Hearing screen • Otosopic exam
Sensory	<ul style="list-style-type: none"> • Sharp • Light touch • Proprioception • Vibration
Motor	<ul style="list-style-type: none"> • Power • Coordination
Vestibular	<ul style="list-style-type: none"> • Dynamic activity • Positional testing
Functional Activities	<p><u>Sitting and standing</u></p> <ul style="list-style-type: none"> • Romberg with eyes open/closed • Single leg stance <p><u>Balance</u></p> <p><u>Transfers</u></p> <ul style="list-style-type: none"> • Supine ↔ sit • Sit ↔ stand <p><u>Gait</u></p> <ul style="list-style-type: none"> • Walking • Tandem walking • Turning

Figure 10.1 Clinical Assessment of Balance

Instructions	<p><i>The 10 Second Balance Screen:</i></p> <p>Age 49 and Under: Ask the subject to stand on one leg, arms free to move. He or she can choose which leg to stand on and are allowed to alternate between legs in between trials. Patients perform the tests with eyes closed. A subject who requests help to assume a testing position is allowed to use the investigator's arm to steady him- or herself prior to starting the timed trials. No instructions are given regarding the subject's knee position. Timing starts when the subject assumes the proper position and indicates that he or she is ready to begin the test. Timing stops when the subject disengages from the starting position or reaches the 30-second time limit. The best of three trials is taken for the result.</p> <p>Age 69 and Under: Ask the subject to stand with one foot just in front of the other with arms free to move (Tandem Romberg). He or she can choose which leg to be in front and can change position in between trials. Patients perform the tests with eyes closed. A subject who requests help to assume a testing position is allowed to use the investigator's arm to steady him- or herself prior to starting the timed trials. Timing starts when the subject assumes the proper position and indicates that he or she is ready to begin the test. Timing stops when the subject disengages from the starting position or reaches the 30-second time limit. The best of three trials is taken for the result.</p> <p>Age 70 and Older: Ask the subject to stand on one leg, arms free to move. He or she can choose which leg to stand on and is allowed to alternate between legs in between trials. Patients perform the tests with eyes open. A subject who requests help to assume a testing position is allowed to use the investigator's arm to steady him- or herself prior to starting the timed trials. No instructions are given regarding the subject's knee position or visual fixation. Timing starts when the subject assumes the proper position and indicates that he or she is ready to begin the test. Timing stops when the subject disengages from the starting position or reaches the 30-second time limit. The best of three trials is taken for the result.</p> <p>Any test score of 10 seconds or less suggests balance impairment.</p>										
	One leg standing (eyes open)										
	Normative Data	Decade	Mean	SD	Median	Perc 05	Interquartile range	Perc 95	Valid N	% 30 s	% 10 s
		3	30.00	.00	30.00	30.00	30.00 - 30.00	30.00	N = 74	100	100
	4	30.00	.00	30.00	30.00	30.00 - 30.00	30.00	N = 43	100	100	
	5	29.64	2.06	30.00	25.91	30.00 - 30.00	30.00	N = 32	97	100	
	6	30.00	.00	30.00	30.00	30.00 - 30.00	30.00	N = 30	100	100	
	7	27.74	5.25	30.00	11.59	30.00 - 30.00	30.00	N = 56	80	95	
	8	21.43	10.08	26.33	2.05	13.04 - 30.00	30.00	N = 56	48	86	
One leg standing (eyes closed)											
	3	27.52	6.45	30.00	9.45	30.00 - 30.00	30.00	N = 74	86	96	
	4	27.48	6.48	30.00	8.46	30.00 - 30.00	30.00	N = 43	86	95	
	5	21.77	9.09	24.75	3.94	10.90 - 30.00	30.00	N = 31	45	90	
	6	19.92	9.81	20.90	3.78	10.55 - 30.00	30.00	N = 29	38	79	
	7	8.93	7.54	5.66	1.61	3.32 - 12.13	28.33	N = 56	4	34	
	8	4.87	3.46	3.93	1.18	2.87 - 6.03	11.78	N = 56	0	5	
Tandem Romberg (eyes closed)											
	3	29.94	.43	30.00	30.00	30.00 - 30.00	30.00	N = 58	98	100	
	4	30.00	.00	30.00	30.00	30.00 - 30.00	30.00	N = 42	100	100	
	5	28.82	4.66	30.00	11.46	30.00 - 30.00	30.00	N = 32	94	97	
	6	28.03	4.87	30.00	13.57	29.70 - 30.00	30.00	N = 28	82	100	
	7	17.96	10.33	16.50	4.18	7.66 - 30.00	30.00	N = 56	36	64	
	8	13.20	9.50	11.26	2.27	4.68 - 18.74	30.00	N = 56	16	54	
Cut-Off	It is recommended that a 10-second time limit per decade is used to delineate poor performance.										

Content based on Vereek, Wuyts, Truijen & Van de Heyning (2008) with normative data tables taken from the paper. © 2008 Informa Healthcare, International Journal of Audiology (<http://informahealthcare.com/ijoa>). Reproduced with permission.

RECOMMENDATIONS FOR PERSISTENT VESTIBULAR (BALANCE/DIZZINESS) DYSFUNCTION

		GRADE
10.1	Evaluation should include a thorough neurologic examination that emphasizes vision, vestibular, balance and coordination, and hearing. See <u>Table 10.1</u> for specific exam details. ^a	C
10.2	If symptoms of benign positional vertigo are present, the Dix-Hallpike manoeuvre (see <u>Appendix 10.1</u>) should be used for assessment.	A
10.3	A canalith repositioning manoeuvre (<u>Appendix 10.1</u>) should be used to treat benign positional vertigo if the Dix-Hallpike manoeuvre is positive.	A
10.4	For persons with functional balance impairments and screening positive on a balance measure, consideration for further balance assessment and treatment by a qualified health care professional may be warranted pending clinical course.	C
10.5	Vestibular rehabilitation therapy is recommended for unilateral peripheral vestibular dysfunction.	A

a. Adapted from the VA/DoD Management of Concussion/Mild Traumatic Brain Injury Clinical Practice Guideline (VA/DoD, 2009).

RECOMMENDATIONS FOR PERSISTENT VESTIBULAR (BALANCE/DIZZINESS) DYSFUNCTION (CONTINUED)		GRADE
10.6	When the patient identifies a problem with hearing the following steps should be followed: 1. Perform an otologic examination. 2. Review medications for ototoxicity. 3. Refer to audiology for hearing assessment if no other apparent cause is found. ^a	C
10.7	When the patient identifies a problem with nausea the following steps should be followed: 1. Define triggers and patterns of nausea and conduct appropriate investigations as required. 2. Assess medication list for agents that may cause or worsen GI symptoms. 3. Perform oropharyngeal examination. 4. Assess vision and vestibular/balance systems. ^a	C

Vision Dysfunction

Patients presenting with vision disorders post-TBI may display anomalies of visual acuity, accommodation, version movements, vergence movements, photosensitivity, visual field integrity and ocular health—collectively termed post-trauma vision syndrome (PTVS; [Table 10.2](#)).⁷⁻⁹ Practitioners should take a detailed history of any persistent vision symptoms and perform examinations to detect potentially unrecognized visual deficits or take note of the specific type of visual disorder the patient is experiencing.^{8,10} Mild TBI patients with advanced ocular health changes and complex strabismic anomalies should be referred to a neuro-ophthalmologist.¹¹⁻¹³ Otherwise, patients who experience changes in accommodation, version or vergence movements, photosensitivity, and visual field integrity are amenable to rehabilitative techniques rendered by qualified optometrists.^{8,10,11} See [Table 10.3](#).

There is some current evidence that optometric vision rehabilitation can be an important modality in the rehabilitation of these patients in certain situations.^{7,8,10,14} It should therefore be offered as a possible option for the treatment and management of persistent vision disorders. Treatment may include rehabilitative interventions such as vision therapy, reading spectacles, prism spectacles, and/or tinted spectacles.^{8,12,14}

Table 10.2 Post-Trauma Vision Syndrome (PTVS) Definitions

Accommodation: The ability to clearly focus the lens of the eye for clear near vision. This ability is gradually lost with increased age (45 yrs +) as a result of loss of elasticity of the lens and its surrounding muscles.
Version Movements: The movement of both eyes in the same direction – easily tested by following a near target in an “H” pattern about 40 cm from the patient.
Vergence Movements: Convergence and divergence eye movements, which enable accurate depth perception. Supra- and infra-vergence relate to the vertical fusional movements of the eyes.

Table 10.3 Common Visual Symptoms and Associated Visual Deficits

Symptom	Possible Visual Deficit
Blurry vision	Accommodative dysfunction
Reading comprehension or efficiency problems	Version eye movement deficits or visual perceptual processing deficits
Diplopia	Vergence eye movement deficits
Eyestrain/headaches	Accommodative or vergence dysfunction
Sensitivity to light/glare	Abnormal light-dark adaptation, photosensitivity
Dizziness	Impaired vestibular-ocular reflex and motion perception
Spatial deficits	Impaired visual field or visual processing deficits

a. Adapted from the VA/DoD Management of Concussion/Mild Traumatic Brain Injury Clinical Practice Guideline (VA/DoD, 2009).

RECOMMENDATIONS FOR PERSISTENT VISION DYSFUNCTION		
		GRADE
10.8	Take an appropriate case history, including questions on visual blur, scanning/reading ability, light sensitivity, diplopia, eyestrain, motion sensitivity, and spatial deficits (indicating loss of visual field integrity). See Table 10.2 for a detailed description of symptoms and their related vision dysfunction.	C
10.9	Perform tests of visual acuity, extra-ocular motility, vergence, visual fields, pupils, and fundoscopy. See Appendix 10.2 for an explanation of screening techniques.	C
10.10	Other functional vision changes should be given consideration for referral to a qualified optometrist specializing in neuro-optometric rehabilitation for vision therapy.	B

RESOURCES	
APPENDICES	
1	Dix-Hallpike Manoeuvre and Particle Repositioning Manoeuvre (PRM) Appendix 10.1
2	Screening Techniques for Vision Dysfunction Appendix 10.2
TABLES	
1	Important Components to Include in the Neurological Exam Table 10.1
2	Post-Trauma Vision Syndrome (PTVS) Definitions Table 10.2
3	Common Visual Symptoms and Associated Visual Deficits Table 10.3
FIGURES	
1	Clinical Assessment of Balance Figure 10.1

References

- Hillier SL, Hollohan V. Vestibular Rehabilitation for unilateral peripheral vestibular dysfunction. Cochrane Database of Systematic Reviews Library. 2007; Issue 4, Art. No.: CD005397. DOI:10.1002/14651858.CD005397.pub2.
- Maskell F, Chiarelli P, Isles R. Dizziness after traumatic brain injury: overview and measurement in the clinical setting. *Brain Injury*. 2006; 20:293-305.
- Parnes LS, Agrawal SK, Atlas J. Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *Canadian Medical Association Journal*. 2003;169:681-693.
- Iverson GL, Kaarto ML, Koehle MS. Normative data for the balance error scoring system: implications for brain injury evaluations. *Brain Injury*. 2008;22:147-152.
- Vereeck L, Wuyts F, Truijen S, Van de Heyning P. Clinical assessment of balance: normative data, and gender and age effects. *International Journal of Audiology*. 2008;47:67-75.
- Hansson EE, Mansson NO & Hakansson A. Effects of specific rehabilitation for dizziness among patients in primary health care. A randomized controlled trial. *Clinical Rehabilitation*. 2004;18:558-565.
- Green W, Ciuffreda KJ, Thiagarajam P, Szymanowicz D, Ludlam DP, Kapoor N. Accommodation in mild traumatic brain injury. *Journal of Rehabilitative Research & Development*. 2010;47(3):183-200.
- Fox RS. The rehabilitation of vergence and accommodative dysfunctions in traumatic brain injury. *Brain Injury Professional*. 2005;2(3):12-15.
- Ciuffreda KJ, Rutner, D, Kapoor N, Suchoff IB, Craig S, Han ME. Vision therapy for oculomotor dysfunctions in acquired brain injury: A retrospective analysis. *Optometry*. 2008;79:18-22.
- Cohen AH. The role of optometry in the management of vestibular disorders. *Brain Injury Professional*. 2005;2(3):8-10.
- Rutner D, Kapoor N, Ciuffreda KJ, Craig S, Han ME, Suchoff IB. Occurrence of ocular disease in traumatic brain injury in a selected sample: A retrospective analysis. *Brain Injury*. 2006;20(10):1079-1086.
- Hillier CG. Vision rehabilitation following acquired brain injury: A case series. *Brain Injury Professional*. 2005;2(3):30-32.
- Kapoor N, Ciuffreda KJ. Vision disturbances following traumatic brain injury. *Current Treatment Options in Neurology*. 2002;4:271-280.
- Ciuffreda KJ, Suchoff IB, Marrone MA, Ahman E. Oculomotor rehabilitation in traumatic brain-injured patients. *Journal of Behavioural Optometry*. 1995;7:31-38.

Appendix 10.1

Dix-Hallpike Manoeuvre and Particle Repositioning Manoeuvre (PRM)*




Fig. 6: Dix-Hallpike manoeuvre (right ear). The patient is seated and positioned so that the patient's head will extend over the top edge of the table when supine. The head is turned 45° toward the ear being tested (position A). The patient is quickly lowered into the supine position with the head extending about 30° below the horizontal (position B). The patient's head is held in this position and the examiner observes the patient's eyes for nystagmus. In this case with the right side being tested, the physician should expect to see a fast-phase counter-clockwise nystagmus. To complete the manoeuvre, the patient is returned to the seated position (position A) and the eyes are observed for reversal nystagmus, in this case a fast-phase clockwise nystagmus.




Fig. 7: Liberatory manoeuvre of Semont (right ear). The top panel shows the effect of the manoeuvre on the labyrinth as viewed from the front and the induced movement of the canaliths (from blue to black). This manoeuvre relies on inertia, so that the transition from position 2 to 3 must be made very quickly.

* Taken from Parnes LS, Agrawal SK, Atlas J. Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *Canadian Medical Association Journal*. 2003;169:681-693. For links to video demonstrations of the above manoeuvres, please see [Appendix B](#).

Appendix 10.2

Screening Techniques for Vision Dysfunction

Visual Acuity	Visual acuity should be performed at both distance and near with each eye, with their current prescription (if applicable).
Extra-ocular Motility	The “Broad H” Test is designed to assess the action of all 6 extraocular muscles around each eye. Have the patient follow a penlight as it is moved into the patient’s right and left field, as well as upwards and downwards in both right and left gaze, making a large “H” pattern out to at least 30-40 degrees (shoulder width as a rule of thumb). The movements should be full and smooth, without diplopia or eyestrain.
Vergence	The ability for the eyes to converge as a team should also be assessed via the Near Point of Convergence test. As a penlight is slowly brought inward towards the patient’s nose, the patient is asked to report when the light “breaks into two” (diplopia). The normal point of convergence is approximately 8cm or less from the nose. If one eye turns outwards, or the patient report diplopia is greater than 8 cm, further investigation is warranted.
Pupils	Pupils should be equal, round and reactive to light without afferent pupillary defect.
Fundoscopy	The internal retinal examination should reveal healthy, distinct optic nerves, maculae and retinal tissue.

Appendix A

Project Members

PROJECT TEAM MEMBERS

Project Team Leader

Shawn Marshall, MD, MSc, FRCPC
Ottawa Hospital Research Institute

Project Investigators

Mark Bayley, MD, FRCPC
Toronto Rehabilitation Institute, University Health Network

Lindsay Berrigan, PhD
Dalhousie University

Scott McCullagh, MD, FRCPC
Sunnybrook Health Science Center

Donna Ouchterlony MD, CCFP
St. Michael's Hospital

Diana Velikonja, PhD, CPsych
Hamilton Health Sciences

Project Coordinator

Kelly Weegar, PhD (Candidate)
Ottawa Hospital Research Institute

Student Research Assistants

Andrew Bayne, BSc (Candidate)
McGill University

Natalie Fersht, BSocSc (Candidate)
University of Ottawa

Hayley MacLeay, BA & Sc (Candidate)
McGill University

Chantal Rockwell, BA (Hons)
Ottawa Hospital Research Institute

Kelsey Scheier, BSc (Candidate)
University of Waterloo

EXPERT CONSENSUS GROUP

Robert Brison MD, MPH, FRCP
Queen's University

Carol Cancelliere DC, MPH
Toronto Western Research Institute, University Health Network

Angela Colantonio PhD, OT
University of Toronto, Toronto Rehabilitation Institute

Victor Coronado, MD, MPH
*Centers for Disease Control and Prevention (USA)**

Nora Cullen, MD, MSc, FRCPC
Toronto Rehabilitation Institute, University Health Network

Lisa Fischer MD, CCFP, DipSportMed
Western University

Anne Forrest, PhD
mTBI Survivor

Bryan Garber, MD
National Defence, Government of Canada

Jonathan Gladstone, MD, FRCPC
University of Toronto

Wayne Gordon, PhD, ABPP/Cn
Mount Sinai School of Medicine

Robin Green, PhD, CPsych
Toronto Rehabilitation Institute, University Health Network

Grant Iverson, PhD, RPsych
University of British Columbia

Corinne Kagan BA, BPS Cert
Ontario Neurotrauma Foundation

Vicki Kristman, PhD
Lakehead University

Andrea Laborde, MD
Royal Hobart Hospital (Australia)

* The recommendations in this document are those of the Ontario Neurotrauma Foundation, identified by the guideline development team and expert consensus group members, and do not necessarily represent agreement of or endorsement by the Centers for Disease Control and Prevention.

Patricia McCord, DC

Willie Miller, MD, FRCPC
University of Ottawa

Gregory O'Shanick, MD
Center for Neurorehabilitation Services (USA)

Marie-Christine Ouellet, PhD
Université Laval

Angela Peddle, OD, FCOVD
University of Waterloo

Laura Rees, PhD, CPsych
The Ottawa Hospital Rehabilitation Center

Wendy Shama MSW, RSW
mTBI Survivor

Mary Stergiou-Kita, PhD
University of Toronto

Rob van Reekum, MD, FRCPC
University of Toronto

Catherine Wiseman-Hakes, PhD, Reg CASLPO
Hôpital du Sacré-Coeur de Montréal

Mark Rapoport, MD, FRCPC
Sunnybrook Health Sciences Center, University of Toronto

Noah Silverberg, PhD, RPsych, ABPP
University of British Columbia

Bonnie Swaine, PT, PhD
Université de Montréal

Charles Tator, PhD, MD
University of Toronto

STAKEHOLDERS/AUDITORS

Humberto Laranjo
Emergency Nurses Association of Ontario

Ruth Wilcock
Ontario Brain Injury Association

Roger Zemek
Children's Hospital of Eastern Ontario

Injury Claims Managers from the Property and
Casualty Insurance Industry

EXTERNAL REVIEWERS

Markus Besemann, LCol, BSc, MD, FRCP, DipSportMed
Canadian Forces Health Services - Rehabilitation Medicine

Pierre Côté, DC, PhD
University of Ontario Institute of Technology

David Cifu, MD
*Virginia Commonwealth University, US Department of
Veterans Affairs*

Appendix B

Other Links/References for Resources to Consider

Section 10: Persistent Vision & Vestibular (Balance/Dizziness) Dysfunction

Balance Error Scoring System (BESS)

A portable and objective method of assessing static postural stability. More specifically, the BESS can be used to assess the effects of traumatic brain injury on static postural stability. The BESS utilizes a combination of stances (feet in a narrow stance, preferably touching; single leg stance; and tandem stance) and footing surfaces (bare feet on the floor or a medium density foam surface).

Guskiewicz KM. Postural stability assessment following concussion: one piece of the puzzle. *Clinical Journal of Sports Medicine*. 2001;11:182–189.

Links for Dix-Hallpike & Repositioning Manoeuvre Video Demonstrations

<http://www.youtube.com/watch?v=kEM9p4EX1jk>

<http://www.youtube.com/watch?v=1-hsUU7MDqc>

<http://www.youtube.com/watch?v=RQV-oz0baFM>